

Letter to the editor: Further identification of a measles variant displaying mutations impacting molecular diagnostics, Northern Italy, 2024

Clara Fappani^{1,2,3,4}, Maria Gori^{1,3,4}, Silvia Bianchi^{1,3}, Marta Canuti⁵, Daniela Colzani^{1,3}, Melissa Baggieri⁶, Silvia Gioacchini⁶, Emilio D'Ugo⁶, Elisabetta Tanzi^{1,3}, Fabio Magurano⁶, Antonella Amendola^{1,3}

1. Department of Health Sciences, Università degli Studi di Milano, Milan, Italy
2. Department of Clinical Sciences and Community Health, Università degli Studi di Milano, Milan, Italy
3. Coordinated Research Center "EpiSoMI", Università Degli Studi di Milano, Milan, Italy
4. These authors contributed equally to this work and share first authorship.
5. Department of Veterinary and Animal Sciences, University of Copenhagen, Frederiksberg, Denmark
6. Department of Infectious Diseases, Istituto Superiore di Sanità, Rome, Italy

Correspondence: Antonella Amendola (antonella.amendola@unimi.it)

Citation style for this article:

Fappani Clara, Gori Maria, Bianchi Silvia, Canuti Marta, Colzani Daniela, Baggieri Melissa, Gioacchini Silvia, D'Ugo Emilio, Tanzi Elisabetta, Magurano Fabio, Amendola Antonella. Letter to the editor: Further identification of a measles variant displaying mutations impacting molecular diagnostics, Northern Italy, 2024. *Euro Surveill.* 2024;29(7):pii=2400079. <https://doi.org/10.2807/1560-7917.ES.2023.29.7.2400079>

Article submitted on 07 Feb 2024 / accepted on 14 Feb 2024 / published on 15 Feb 2024

To the editor: We read with great interest the article by Pérez-Rodríguez et al. [1], reporting the circulation of measles virus (MeV) variants (genotype D8) displaying three mutations impacting molecular diagnostics. The mutations occurred in the 450nt-long C-terminal (N-450) region of the nucleoprotein (N) gene that is usually the target of real-time reverse-transcription (RT)-PCR assays commonly employed by surveillance laboratories. In detail, the three T-to-C synonymous substitutions were located within the annealing site of the reverse primer recommended by the United States Centers for Diseases Control and Prevention (CDC, Atlanta) [2]. This results in a slight loss of test sensitivity.

As a Subnational Reference Laboratory of the measles and rubella surveillance network MoRoNet [3], we currently employ a real-time RT-PCR assay using the primers and probe described by Hübschen et al. [4], to amplify a fragment of 114nt of the N gene that does not fall within the N-450 region. Since January 2024, we have confirmed five measles cases in the surveyed area, which includes the Metropolitan City of Milan and surrounding areas in Lombardy, Northern Italy (ca 4 million inhabitants). This region borders Switzerland, where the study by Pérez-Rodríguez and colleagues was conducted. The five Lombard cases were all classified as genotype D8 and determined to be sporadic since we found no clear epidemiological link among them. In three of these cases, a recent history of travel was reported, and destinations included Uzbekistan, Thailand, and Southern Italy. In particular, two of these cases, each with a travel history to either Southern Italy or Thailand, were infected with MeV strains characterised by the three mismatches described by Pérez-Rodríguez et al. [1].

Both cases were confirmed through the detection of virus-specific IgM in serum with an enzyme-linked immunosorbent assay, and of MeV-RNA in both urine and oropharyngeal swab samples by real-time RT-PCR [4]. MeV strains were genotyped by sequencing the N-450 region as recommended by the World Health Organization (WHO) [5,6]. Both sequences were annotated and deposited in the WHO Measles Virus Nt Surveillance (MeaNS2) database [7] with the distinct sequence identifier (DSId) 8491, and in GenBank [8] (accession numbers: PP334141–PP334142). Notably, the DSId was different from the one (i.e. 8248) detected by Pérez-Rodríguez et al. [1]. Basic Local Alignment Search Tool (BLAST) analysis showed that they shared 99.7% identity with strains identified in Moscow (Russia) in 2023 (GenBank accession numbers: OR290099–OR290102, OR840959, OR840960) [9].

None of the other 614 strains (453 D8 and 161 B3) detected by our laboratory between 2017 (the beginning of surveillance activities) and 2023 possessed these mutations. However, our results suggest that MeV with the specific mutations detected through the Swiss molecular surveillance are already circulating in Italy, in line with the results obtained by Pérez-Rodríguez et al., who reported a local case with travel history to Italy.

We commend Pérez-Rodríguez and colleagues for the rapid communication as they raise awareness on the circulation of a MeV variant that can be detected with reduced sensitivity by many currently used diagnostic tests. This letter confirms their finding and reports that the identified variant is spreading. This highlights the

importance of promptly updating diagnostic tests to detect all currently circulating MeV strains.

This article is copyright of the authors or their affiliated institutions, 2024.

Acknowledgements

The authors wish to thank Sabrina Senatore and the staff of the local health authorities of Milan involved in measles case investigations, and the Directorate General of Welfare of the Lombardy Region for the help in retrieving the epidemiological data from the regional database SMI.

Conflict of interest

None declared.

Authors' contributions

Conceptualisation: AA; drafting of the letter: CF, MG, and SB; contributed analysis: CF, MG, SB, DC, MB, SG, and ED; critical revision of the letter: MC, ET, FM, and AA.

References

1. Pérez-Rodríguez FJ, Cherpillod P, Thomasson V, Vetter P, Schibler M. Identification of a measles variant displaying mutations impacting molecular diagnostics, Geneva, Switzerland, 2023. *Euro Surveill.* 2024;29(5):2400034. <https://doi.org/10.2807/1560-7917.ES.2024.29.5.2400034> PMID: 38304951
2. Hummel KB, Lowe L, Bellini WJ, Rota PA. Development of quantitative gene-specific real-time RT-PCR assays for the detection of measles virus in clinical specimens. *J Virol Methods.* 2006;132(1-2):166-73. <https://doi.org/10.1016/j.jviromet.2005.10.006> PMID: 16274752
3. Istituto Superiore di Sanità. Moronet Lab | liberi dal morbillo e dalla rosolia [Moronet; free from measles and rubella]. Available from: <https://moronetlab.it/>
4. Hübschen JM, Kremer JR, De Landtsheer S, Muller CP. A multiplex TaqMan PCR assay for the detection of measles and rubella virus. *J Virol Methods.* 2008;149(2):246-50. <https://doi.org/10.1016/j.jviromet.2008.01.032> PMID: 18353451
5. World Health Organization (WHO). Manual for the laboratory diagnosis of measles and rubella virus infection, 2nd edition. Geneva: WHO; 2018. Available at: <https://www.who.int/publications-detail-redirect/WHO-IVB-07.01>
6. Bianchi S, Frati ER, Lai A, Colzani D, Ciceri G, Baggieri M, et al. Genetic characterisation of Measles virus variants identified during a large epidemic in Milan, Italy, March-December 2017. *Epidemiol Infect.* 2019;147:e80. <https://doi.org/10.1017/S0950268818003606> PMID: 30869017
7. Williams D, Penedos A, Bankamp B, Anderson R, Hübschen J, Ben Mamou M, et al. Update: circulation of active genotypes of measles virus and recommendations for use of sequence analysis to monitor viral transmission. *Wkly Epidemiol Rec.* 2022;39(97):481-92.
8. Benson DA, Cavanaugh M, Clark K, Karsch-Mizrachi I, Lipman DJ, Ostell J, et al. GenBank. *Nucleic Acids Res.* 2013;41(Database issue):D36-42. PMID: 23193287
9. BLAST. Basic Local Alignment Search Tool. Available at: <https://blast.ncbi.nlm.nih.gov/Blast.cgi>

License, supplementary material and copyright

This is an open-access article distributed under the terms of the Creative Commons Attribution (CC BY 4.0) Licence. You may share and adapt the material, but must give appropriate credit to the source, provide a link to the licence and indicate if changes were made.

Any supplementary material referenced in the article can be found in the online version.